A SEARCH FOR CANDIDATE RADIO SUPERNOVA REMNANTS IN THE NEARBY IRREGULAR STARBURST GALAXIES NGC 4214 AND NGC 4395 (Serb. Astron. J. № 170 (2005), 101)

B. Vukotić¹, I. Bojičić¹, T. G. Pannuti² and D. Urošević¹

¹Department of Astronomy, Faculty of Mathematics, University of Belgrade Studentski trg 16, 11000 Belgrade, Serbia and Montenegro

²Spitzer Science Center, California Institute of Technology, MS 220-6, Pasadena, CA 91125, USA; tpannuti@ipac.caltech.edu

In the paper published in Serb. Astron. J. No. 170 we made a mistake in Eq. 5. It should read:

$$B_{\rm min} = (4.5)^{2/7} (1+\xi)^{2/7} c_{12}^{2/7} f^{-2/7} (d/2)^{-6/7} L_{\rm radio}^{2/7}$$

The mistake was also made in calculations obtained with Eq. 5. Column 3 of Table 4 should read:

| Candidate Radio SNR | B_{\min} (mG) |
|------------------------|-----------------|
| α | 0.29 |
| β | 0.28 |

Row 6, column 4 of Table 5 should read:

| Galaxy | $\begin{array}{c} \text{Mean} \\ B_{\min} \\ \text{(mG)} \end{array}$ |
|----------|---|
| NGC 4214 | 0.29 |