

## STARK BROADENING OF Be II SPECTRAL LINES

M. S. DIMITRIJEVIĆ<sup>1,2</sup>, M. CHRISTOVA<sup>3</sup> AND S. SAHAL-BRÉCHOT<sup>2</sup>

<sup>1</sup>*Astronomical Observatory, Volgina 7, 11060 Belgrade 38, Serbia*

<sup>2</sup>*Laboratoire d'Etude du Rayonnement et de la Matière en Astrophysique,  
Observatoire de Paris, UMR CNRS 8112, UPMC, 5 Place Jules Janssen,  
92195 Meudon Cedex, France*

<sup>3</sup>*Department of Applied Physics, Technical University-Sofia, 1000 Sofia,  
Bulgaria*

E-mail: mchristo@tu-sofia.bg

Calculations of the Stark broadening parameters of singly ionized beryllium spectral lines will be presented. The work is based on the semi-classical theory developed in (Sahal-Bréchet, 1969a,b). Values of energy levels in (Kramida *et al.* 2005) are used.

### References

- Sahal-Bréchet, S.: 1969a, *A&A*, **1**, 91.  
Sahal-Bréchet, S.: 1969b, *A&A*, **2**, 322.  
Kramida, A. E. and Ryabtsev, A. N.: 2005, *Phys. Scr.* **72**, 309.